

EFFECT OF IONIZING (GAMMA) AND NON-IONIZING (UV) RADIATION ON THE DEVELOPMENT OF *TRICHOGRAMMA EUPROCTIDIS* (HYMENOPTERA: TRICHOGRAMMATIDAE)

AYDIN S. TUNCBILEK¹, FAHRIYE SUMER ERCAN² and ULKU CANPOLAT¹

¹Department of Biology, Faculty of Sciences, Erciyes University, 39039 Kayseri, Turkey

²Department of Biology, Faculty of Sciences and Literature, Bozok University, Yozgat, Turkey

Abstract - The potential of using gamma and ultraviolet radiation as an alternative treatment to increase the efficiency of *Trichogramma euproctidis* (Girault 1911) (Hymenoptera: Trichogrammatidae) was investigated in the laboratory. The developmental and adult stages of *T. euproctidis* were exposed to gamma radiation of different doses (0-30 Gy) and ultraviolet radiation of 254 nm wavelengths (UV-C) for different durations (0-10 min) to assess their effect on each of the instars and their potential in breaking the developmental cycle of the egg parasitoid. The LD₅₀ values for eggs, prepupae, pupae and adults were 8.1, 10.0, 22.7 and 9.5 Gy for gamma radiation and 9.5, 0.12, 2.0 and 11.9 min for UV radiation, respectively. The pupa and adult stages were more radioresistant to both gamma and UV radiation. The most interesting and unexpected result obtained for the prepupal stage was that UV radiation has a greater effect on prepupal stages than gamma radiation.

Key words: Parasitoid, *Trichogramma euproctidis*, longevity, gamma radiation, UV radiation, LD₅₀ and LD₉₉

INTRODUCTION

Nuclear techniques could play a vital role in enabling the cost-effective mass production of beneficial insects for use in augmentative biological control. There are significant opportunities for increasing the use of classical and augmentative biological control through nuclear techniques for the production and release of biological control agents (Brower, 1982; Greany, 1999).

The Ultraviolet (UV) portion of the spectrum has been widely used as a germicide and as an attractant for insects (Bruce, 1975) in embryological-physiological studies for the surface disinfection of insect eggs from pathogens (Bodenstein, 1953; Guerra et al., 1968) and for the suppression of insects and different stages of the life cycle (Beard, 1972; Baden

et al., 1996). Among these, UV-B/C radiation is very important in influencing biological systems (Kovacs and Keresztes, 2002). Egg parasitoids in the genus *Trichogramma* are one of the most important biological control agents of pest insects in different crops. Augmentative releases of the *Trichogramma* species require mass rearing. The eggs of *Ephesttia kuehniella* Zell. (Lepidoptera: Pyralidae) are used as an alternative host for rearing *Trichogramma*. *E. kuehniella* larvae that hatch from unparasitized eggs and can attack parasitized ones. To prevent host larvae hatch, ultra violet light (UV) can be used to kill host eggs. Gamma radiation is an ionizing radiation type and has a very high penetrating ability (Ayvaz and Tunçbilek, 2006). Nevertheless, host eggs exposed to gamma irradiation can be used as a food for *Trichogramma* rearing (Mansour, 2010). In Turkey, some studies have been conducted on the effect of gamma

rays on *Trichogramma* (Tuncbilek and Canpolat, 2003; Tuncbilek and Ayvaz, 2001). Unfortunately, no research has been conducted on the effect of ultraviolet radiation on the *Trichogramma*. Keeping in mind the importance and feasibility of the use of UV-rays, the present investigation was undertaken to evaluate the effect of gamma and UV-irradiation on the commercially relevant aspects of *Trichogramma* rearing. It is, however, difficult to make direct comparisons between gamma and UV radiation studies as the level of gamma and UV dose achieved is not always stated, and radiation intensities vary with sources. The ability of UV to penetrate surfaces is very limited, and thus its effect is limited to surfaces, whereas gamma rays can penetrate deep into tissues. In principle, ultraviolet C (UVC) radiation may provide an effective means of combating pest infestations associated with the structure of a building and may serve as a potential new hygiene measure. Although the use of gamma radiation as a method of pest control has been extensively investigated (Tuncbilek, 1995; Tuncbilek and Kansu, 1996; Tuncbilek, 1997; Carpenter et al., 2001; Makee and Saour, 2003; Bloem et al., 2006), the use of UV-C radiation has not been widely studied due to the lack of penetration through substrates (Bruce and Lum, 1978).

These results led us to explore the idea that radiation is very important in enhancing parasitoid effectiveness in biological systems. The research described herein was conducted to examine the possible effects of UV and gamma radiation use in *T. euproctidis*.

MATERIALS AND METHODS

Rearing of Ephestia kuehniella

All insects were taken from an *E. kuehniella* stock culture kept in the laboratory in 3 l plastic jars covered with mesh. The larvae were reared on a pesticide-free mixture consisting of 1 kg wheat flour, 5% yeast and 30 g of wheat germ. Throughout the rearing, cultures were kept in a rearing room equipped with a controlling system, at $27\pm 1^\circ\text{C}$ and $70\pm 5\%$ R.H., 16:8 h L:D. Light was controlled using a 24 h time switch and fluorescent tubes.

Rearing of Trichogramma euproctidis

The wasp *Trichogramma euproctidis* (Girault 1911) was reared on the eggs of the flour moth and kept in test tubes at $27\pm 1^\circ\text{C}$, 70–80% RH, 16:8 h L:D. Strips of lightweight cardboard (2.5 by 4 cm) were brushed with gum arabica. The eggs were sprinkled onto these cards and placed in tubes along with adult *T. euproctidis*. Individual *T. euproctidis* females (24 h old) were prepared for the tests by isolating them in small tubes. To do this, the adults were scattered from the rearing tubes onto white paper and captured by placing test tubes (180 mm long and 18 mm in diameter) over them. When the parasites walked up into the tube towards the source of light, their sex could be easily determined under a binocular microscope and then the egg card was placed into the tube (Wührer and Hassan, 1993). After 24 h, the egg cards were removed from the tubes and incubated in controlled conditions.

Gamma and UV radiation sources

T. euproctidis eggs, prepupae, pupae and adults were placed in glass tubes and irradiated in a calibrated ^{137}Cs irradiator (IRACEL OK-GK17) with a source strength of ca 111 TBq and a dose rate of ca 3.20 Gy/min. Dose rate was verified with Fricke dosimetry. They were exposed to 7 dose levels ranging from 0, 5, 10, 15, 20, 25 and 30 Gy. An unirradiated control population (0 dose) was similarly started.

All the developmental stages of the wasp were irradiated with UV-rays of 254 nm wavelength at different duration: 0, 2, 4, 6, 8 and 10 min (Mineralight Lamp, shortwave UV, 254 nm 215–250 V, 56–60 Hz, 0.12A). For irradiation, the test insects were kept in 10 cm quartz tubes placed on the surface 5 cm away from the lamp. During irradiation, the room was illuminated with two 40W fluorescent bulbs and the temperature of the laboratory was $27\pm 1^\circ\text{C}$.

Experiments

One-day-old *T. euproctidis* adults were obtained from the stock culture. Male and female adults were

allowed to mate by pairing them inside glass tubes in a growth chamber at $27\pm 1^{\circ}\text{C}$, $70\pm 5\%$ r.h, at a photoperiod of 16:8 h light:dark, respectively. Individual *T. euproctidis* females (24 h old) were prepared for the tests by isolating them in small tubes for laying. Parasitized eggs were isolated into different groups, one for the control, six and five for the different doses of gamma and UV radiation, respectively. The 1-day-old parasitized eggs in each replicate were exposed to gamma radiation by placing the eggs in glass tubes, or UV doses by placing the eggs in quartz tubes for egg stage irradiation of *T. euproctidis*. Each group consisted of five replicates, with 50 eggs/replicate. Control eggs were placed in the room for the designated time without being exposed to radiation. Eggs parasitized by *T. euproctidis* were treated at target gamma radiation doses of 0, 5, 10, 15, 20, 25 and 30 Gy. In the UV radiation test, the eggs were treated at 0, 2, 4, 6, 8 and 10 min exposure. After irradiation, the eggs were returned to their corresponding glass tubes and placed into rearing room.

The prepupae chosen for irradiation were 4-day-old parasitized eggs. This means that the parasitized eggs were irradiated 4 days after laying. The appearance of black deposits on the surface was an indication of the onset of the prepupal stage (Dahlan and Gordh, 1997). Pupae were irradiated 8 days after laying and one-day-old *T. euproctidis* adults were irradiated with the doses mentioned above. After exposure to radiation, irradiated stages and controls were kept in the rearing rooms under the conditions mentioned above. The numbers of parasitized eggs, adult emergence and sex ratio for each stage were scored.

Statistical analysis

Data were subjected to analysis of variance (ANOVA) for the determination of differences between means, and when significant differences occurred, a Tukey-HSD test was applied for mean separation. The data were transformed to square roots before statistical analysis was performed; back-transformed data are presented in the tables and figures. LD_{50} and LD_{99} values were estimated by using probit analysis with doses of radiation as sources of variation (SPSS, 1999).

RESULTS

Egg

The effects of applying different doses of gamma and UV irradiation on the subsequent development of eggs are shown in Fig. 1. It was found that mean parasitization and adult emergence were significantly reduced with both increased gamma radiation doses but not significantly by UV radiation exposure time ($F=22.911$, $df=6$, $P<0.001$; $F=1.568$, $df=5$, $P=0.207$ for gamma and UV radiation, respectively). The effect of gamma radiation was more pronounced than UV radiation at the studied doses. An increase in gamma radiation dose and in time of exposure to UV-rays caused a gradual decrease in the number of hatching eggs. Although very few adults (2.8) emerged from gamma irradiated *T. euproctidis* eggs, more adults (8.6) emerged from UV irradiated eggs after 25 Gy and 10 min, respectively. One-way ANOVA showed that gamma and UV-radiation had significant effects on adult emergence in comparison to controls ($F=35.445$, $df=6$, $P<0.001$; $F=4.879$, $df=5$, $P=0.003$ for gamma and UV radiation, respectively). Gamma radiation was more effective than UV radiation in the egg stage in the studied doses.

Although female emergence was influenced by both gamma and UV radiation ($F=25.666$, $df=6$, $P<0.001$; $F=6.096$, $df=5$, $P=0.001$ for gamma and UV radiation, respectively), male emergence was not changed by the radiation treatments ($F=4.803$, $df=6$, $P=0.002$; $F=1.041$, $df=5$, $P=0.417$ for gamma and UV radiation, respectively).

Prepupae

The results of the effect of gamma and UV radiation on prepupae are shown in Fig. 2. There was a significant reduction in the adult emergence numbers produced by UV-treated females compared to gamma radiation-treated females ($F=60.620$, $df=6$, $P<0.001$; $F=206.324$, $df=5$, $P<0.001$ for gamma and UV radiation, respectively). The effect of UV radiation on adult emergence was more evident than gamma radiation. Although very few adults emerged from both gamma




Fig. 1. Mean (\pm SD) parasitization and adult emergence from gamma and UV irradiated eggs of *T. euproctidis*.




Fig. 2. Mean (\pm SD) adult emergence from gamma and UV irradiated prepupae of *T. euproctidis*.

and UV irradiated *T. euproctidis* prepupae after 25 Gy and 2 min, respectively, no adults developed from prepupae irradiated with higher doses or exposure times. *T. euproctidis* prepupae were unexpectedly more vulnerable to damage from UV radiation than from gamma radiation. To our knowledge, this is the first report of UV radiation being more efficient than gamma radiation on the prepupal stage.

Female and male emergences were influenced by radiation treatments (Female: $F=52.571$, $df=6$,

$P<0.001$; $F=14.378$, $df=5$, $P<0.001$ for gamma and UV radiation, respectively; Male: $F=3.683$, $df=6$, $P=0.008$; $F=7.833$, $df=5$, $P<0.001$ for gamma and UV radiation, respectively).

Pupae

Fig. 3 shows the effect of exposing *T. euproctidis* pupae to gamma and UVC radiation on their subsequent development. It was found that adult emergence decreased with both increased gamma and




Fig. 3. Mean (\pm SD) adult emergence from gamma and UV irradiated pupae of *T. euproctidis*.




Fig. 4. Mean (\pm SD) parasitization and adult emergence from gamma and UV irradiated adults of *T. euproctidis*.

UV radiation doses ($F=11.300$, $df=6$, $P<0.001$; $F=6.793$, $df=5$, $P<0.001$ for gamma and UV radiation, respectively). Adult eclosion from irradiated *T. euproctidis* pupae gradually decreased as the gamma radiation doses increased, but adult eclosion from UV irradiated *T. euproctidis* pupae drastically decreased with duration of exposure to radiation. Although *T. euproctidis* pupae were more tolerant to both UV and gamma radiation, UV radiation was more effective than gamma radiation in the pupal stage.

Gamma and UV radiation did not influence either female emergence ($F=5.190$, $df=6$, $P=0.001$; $F=1.357$, $df=5$, $P=0.275$ for gamma and UV radiation, respectively) or male emergence ($F=1.605$, $df=6$, $P=0.183$; $F=2.479$, $df=5$, $P=0.060$ for gamma and UV radiation, respectively) when irradiated in the pupal stage.

Adult

The results of the effect of gamma and UV exposure




Fig. 5. LD₅₀ values of gamma and UV radiated development stages of *T. euproctidis*.




Fig. 6. Longevity of irradiated adult of *T. euproctidis* with gamma and UV radiation.

on adults are shown in Fig. 4. The mean parasitization ($F=66.586$, $df=6$, $P<0.001$; $F=4.342$, $df=5$, $P=0.060$ for gamma and UV radiation, respectively) and adult emergence ($F=33.970$, $df=6$, $P<0.001$; $F=6.262$, $df=5$, $P=0.001$ for gamma and UV radiation, respectively) decreased with both increased gamma and UV radiation doses. Although *T. euproctidis* adults were more tolerant to both UV and gamma radiation, the effect of gamma radiation on parasitization and adult emergence was more pronounced than UV radiation with the studied doses.

Gamma and UV radiation did not influence either female ($F=53.551$, $df=6$, $P<0.001$; $F=1.616$, $df=5$, $P=0.194$ for gamma and UV radiation, respectively) or male emergence ($F=5.459$, $df=6$, $P=0.001$; $F=2.023$, $df=5$, $P=0.111$ for gamma and UV radiation,

respectively) when UV-treated females were compared to gamma radiation-treated females.

T. euproctidis adults were more resistant to UV radiation than gamma radiation. Only a few adults emerged from irradiated adults after 25 Gy and no adults emerged after 30 Gy of gamma radiation, but adult emergence was observed for all durations of exposure to UV. LD₅₀ values for eggs, prepupae, pupae and adults exposed to gamma radiation were 8.1, 10.0, 22.7 and 9.5 Gy, respectively, and LD₅₀ values for eggs, prepupae, pupae and adults exposed to UV radiation were 9.5, 0.12, 2.0 and 11.9 min, respectively. The comparison of LD₅₀ doses of gamma and UV radiation for the developmental stages and adults of *T. euproctidis* showed that these values differed statistically. The pupae and adult stages were more radi-

oresistant to gamma radiation and to UV radiation, respectively. The egg and prepupal stages were more sensitive to gamma radiation and to UV radiation, respectively.

Longevity

The longevity of *T. euproctidis* adults irradiated with gamma and UV radiation are shown in Fig. 5. Although the longevity of adults irradiated with UV radiation gradually decreased from 13 days to 9 days with increasing doses, it significantly decreased from 11 days to 3 days with increased doses of gamma radiation.

DISCUSSION

In the present study, gamma and ultra-violet radiation (UV) were used for *T. euproctidis* to enhance its parasitization and longevity. A search of databases for research on gamma and UV radiation and *T. euproctidis* did not yield any information on the minimum radiation doses for suppressing/stimulation. Therefore, we chose radiation doses arbitrarily to evaluate the effect of UV doses on the developmental stages of *T. euproctidis*. However, the striking difference in the observed effect could be attributed in part to the difference in the nature of the two types of radiation. The ability of UV to penetrate surfaces is very limited, and thus its effect is limited to surfaces, whereas gamma rays can penetrate deep into tissues. It is difficult to compare the effect of UV irradiation with the effect of gamma irradiation on *T. euproctidis* because of the difficulties involved in relating UV doses to gamma ray doses.

Our study showed that the effect of gamma radiation was more evident than UV radiation in the studied doses. An increase in exposure time to UV-rays caused a gradual decrease in the percentage of hatching eggs. The results obtained in our study may be correlated with the findings of Tilton and Brower (1983), who observed that the embryonic stage of an animal is a time of extreme radiosensitivity and insects are no exception. It was reported that an increase in exposure time to UV-rays caused a gradual

decrease in the percentage of hatching eggs and no hatching occurred after 24 min of exposure in 2- and 3-day-old eggs of *T. confusum* (Faruki et al., 2007). Studies have been made on unlaidd eggs (Hussain et al., 1994), on embryos produced parthenogenetically (Ahmed et al., 1976) and on embryos produced by fertilization (Ghomomu, 1989).

It was observed that the prepupae of *T. euproctidis* were more sensitive to UV rays than the eggs. The prepupae were unexpectedly most vulnerable to damage from UV radiation in comparison to gamma radiation. To our knowledge, this is the first report of UV radiation being more efficient than gamma radiation on the prepupal stage. The results obtained in our study may be correlated with the findings of Daimona et al. (2010) who observed that silkworm prepupae were highly sensitive to UV-B irradiation and their pupation rates dramatically decreased with the increase of radiation dose. Early research into the effects of irradiation on larvae was made using different storage pests. Larvae are more resistant to radiation than eggs and there is a variation in susceptibility between larval stages (Nair and Subramanyam, 1963; Yang and Ducoff, 1971; Allotey, 1985; Johnson and Patrick, 1988; Hasan et al., 1989).

Adult eclosion from irradiated *T. euproctidis* pupae gradually decreased as gamma radiation doses increased, however, adult eclosion from UV irradiated pupae drastically decreased with duration of exposure to radiation. *T. euproctidis* pupae were more tolerant to both UV and gamma radiation. In many ways, the pupal stage is analogous to the embryonic stage when intense cell division and differentiation occur. Therefore, it is not surprising that the overall pattern of pupal radiosensitivity is similar to the pattern exhibited by developing eggs (Tilton and Brower, 1983). Most studies of pupal radiosensitivity have not been precise enough, or at least not correlated with the morphogenetic changes occurring to reveal the fine details in the pattern of radiosensitivity. Irradiation can have several distinct effects on pupae, among which are pupal mortality, adult malformations and delayed adult mortality. The time sequence of mortality in pupae is very difficult to determine

and their failure to emerge is usually used to assess overall pupal mortality. Working with wheat infested by the granary beetle, *Sitophilus granarius* (L.), Cornwell (1966) demonstrated a marked increase in the radiation resistance of the pupae compared with larvae.

Mean parasitization and adult emergence from irradiated females decreased with both increased gamma and UV radiation doses. Also, the effect of gamma radiation on parasitization and adult emergence was more pronounced than UV radiation. Adult insects are more radioresistant than the other stages in the developmental sequence (Tilton and Brower, 1983). In spite of many studies on the effects of radiation on insects, the specific mechanisms whereby the damage is produced and later manifested are still not well understood. Bhuiya et al. (1991) showed a dose-dependent reduction in adult emergence from the irradiated eggs, larvae and pupae of *Sitotroga cerealella* and *Oryzaephilus surinamensis*. Although it is difficult to make direct comparisons between gamma and UV radiation on their biological effects, further research will more precisely identify the wavelengths responsible for prepupal death or the stimulating effect and relationships between gamma and UV radiation in events.

Acknowledgment - The authors would like to thank the Research Fund of Erciyes University for its support through Project Number: FBT-06-84.

REFERENCES

- Ahmed, M.Y., Elbadry, E.A., and Y.S. Salem (1976). Effects of gamma radiation on the reproduction organs of the northern cowpea weevil, *Callosobruchus maculatus* F. *Annales de Zoologie, Ecologie Animale* **8** (2), 279-284.
- Allotey, J. (1985). Study of radiosensitivity of the immature stages of *Corcyra cephalonica* (Stainton) (Lepidoptera: Galleriidae). *Insect Science and its Application* **6** (5), 621-625.
- Ayvaz, A., and A.S. Tunçbilek (2006). Effects of gamma radiation on life stages of the Mediterranean flour moth, *Ephestia kuehniella* Zeller (Lepidoptera: Pyralidae). *Journal of Pest Science*, **79**, 215-222.
- Baden, H.P., Kollias, N., Anderson, R.R., Hopkins, T., and L. Raffery (1996). *Drosophila melanogaster* larvae detect low doses of UVC radiation as manifested by a writhing response. *Archives of Insect Biochemistry and Physiology* **32**, 187-196.
- Beard, R.L. (1972). Lethal action of UV irradiation on insects. *Journal of Economic Entomology* **65**, 650-654.
- Bhuiya, A.D., Ahmed, M., Rezaur, R., Nahar, G., Huda, S.M.S., and S.A.K.M., Hossain (1991). Radiation disinfestation of pulses, oilseeds and tobacco leaves. In insect disinfestation of food and agricultural products by irradiation. *Proceeding of final research co-ordination meeting*, Vienna: IAEA. pp. 27-50.
- Bloem, S., Carpenter, J.E., and S. Dorn (2006). Mobility of mass reared diapaused and nondiapaused *Cydia pomonella* (Lepidoptera: Tortricidae): effect of mating status and treatment with gamma radiation. *Journal of Economic Entomology* **99**, 699-706.
- Bodenstein, D. (1953). Embryonic development. In: Roder KD, editor. *Insect Physiology* pp. 780-822. John Wiley and Sons Publishing.
- Brower, J.H. (1982). Parasitization of irradiated eggs from irradiated adults of the Indian Meal Moth (Lepidoptera: Pyralidae) by *Trichogramma pretiosum* (Hymenoptera: Trichogrammatidae). *Journal of Economic Entomology* **75**, 939-944.
- Bruce, W.A. (1975). Effect of UV-radiation on egg hatch of *Plodia interpunctella* (Lepidoptera: Pyralidae). *Journal of Stored Product Research* **11**, 243-244.
- Bruce, W.A., and P.T.M., Lum (1978). The effects of UV radiation on stored-product insects. In: Caswell GH, Boshoff WH, Daramola AM, Dina SO, Adesuyi SA, Singh SR, Ivbijaro MF, Adeyemi SAO, Taylor TA, editors. *Proceedings of the Second International Working Conference on Stored-Product Entomology*, pp. 271-277. Ibadan, Nigeria.
- Carpenter, J.E., Bloem, K.A., and S. Bloem (2001). Applications of F₁ sterility for research and management of *Cactoblastis cactorum* (Lepidoptera: Pyralidae). *Florida Entomology* **84**, 531-536.
- Cornwell, P.B. (1966). The entomology of radiation disinfestation of grain, Pergamon Press, London. pp. 236.
- Daimon, T, Hirayama, S, Kanai, M, Ruike, Y, Meng, Y, Kosegawa, E, Nakamura, M, Tsujimoto, G., Katsuma, S., and T. Shimada (2010). The silkworm Green b locus encodes a quercetin 5-O-glucosyltransferase that produces green cocoons with UV-shielding properties. *Proceedings of the National Academy of Sciences (PNAS)* **107** (25), 11471-11476.
- Dahlan, A.N., and G. Gordh (1997). Development of *Trichogramma australicum* (Hymenoptera: Trichogrammatidae) at low and high population density in artificial diet. *Entomophaga* **42**, 525-536.

- Faruki, S.I., Das, D.R., Khan, A.R., and M. Khatun (2007). Effects of ultraviolet (254nm) irradiation on egg hatching and adult emergence of the flour beetles, *Tribolium castaneum*, *T. confusum* and the almond moth, *Cadra cautella*. *Journal of Insect Science* 7 (36), 1-6.
- Ghomomu, T.R. (1989). Some observations on the reproduction of adults of the cowpea weevil, *Callosobruchus maculatus* F. resulting from the irradiated stages. *Parasitica* 45 (4), 99-104.
- Greany, P. (1999). The potential role of nuclear techniques in support of the production of biological control agents of insect pests, In: *Use of nuclear techniques in biological control: Managing pests, facilitating trade and protecting environment*. Report of a consultants' group meeting, Vienna, IAEA-314-D4-97CT02288.
- Guerra, A.A., Ouye, M.T., and H.R. Bullock (1968). Effect of ultraviolet irradiation on egg hatch, subsequent larval development, and adult longevity of the tobacco budworm and the bollworm. *Journal of Economical Entomology* 61, 541-542.
- Hasan, M., Khalequzzaman, M, and A.R. Khan (1989). Development of *Tribolium anaphe* irradiated as larvae of various ages with gamma rays. *Entomologia Experimentalis et Applicata* 53, 92-94.
- Hussain, T., Imura, O., and Z.A. Qureshi (1994). Effect of gamma radiation on post-embryonic development following irradiation of *Callosobruchus chinensis* eggs. *Pakistan Journal of Zoology* 26 (1), 7-9.
- Johnson, J.A., and V.V. Patrick (1988). Post-treatment, survival, development, and feeding of irradiated Indian meal moth and Navel orange worm larvae (Lepidoptera: Pyralidae). *Journal of Economical Entomology* 81 (1), 376-380.
- Kovacs, E., and A. Keresztes (2002). Effect of gamma and UV-B/C radiation on plant cells. *Micron* 33, 199-210.
- Makee, H., and G. Saour (2003). Noninherited sterility in irradiated *Phthorimaea operculella* females. *Journal of Applied Entomology* 127, 489-493.
- Mansour, M. (2010). Effects of gamma radiation on the Mediterranean flour moth, *Ephestia kuehniella*, eggs and acceptability of irradiated eggs by *Trichogramma cacoeciae* females. *J. Pest. Sci.* 83, 243-249.
- Nair, K.K., and G. Subramanyam (1963). Effects of variable dose-rate on radiation damage in the rust-red flour beetle, *Tribolium castaneum* Herbst, In: *Radiation and radioisotopes applied to insects of agricultural importance*, pp. 425-429. STI/PUB/74, IAEA, Vienna.
- SPSS. (1999). SPSS Version 10.0. SPSS Inc, 233 S. Wacker Drive, Chicago, Illinois.
- Tilton, E.W., and J.H. Brower (1983). Radiation effects on arthropods. In: Josephson ES, Peterson MS, editors. *Preservation of food by ionizing radiation*, Vol. 2, pp. 269-316. Boca Raton, FL: CRC Press Inc.
- Tuncbilek, A.S., and U. Canpolat (2003). Effects of Stored and Gamma Irradiated Host-Eggs on Some Performance Characteristics of *Trichogramma evanescens*. *European Society of New Methods in Agricultural Research, XXXIIIth Annual Meeting*, Viterbo, Italy p. 96
- Tuncbilek, A.S., and A. Ayvaz (2001). The use of nuclear techniques for the production and colonization of *Trichogramma evanescens* to manage flour moth, *Ephestia kuehniella*, In: *Evaluating the use of nuclear techniques for the colonization and production of natural enemies of agricultural insect pests*, Second Research Co-ordination Meeting Organized by the FAO/IAEA Division of Nuclear Techniques in Food and Agriculture, Tapachula, Mexico, Reproduced by FAO/IAEA Vienna, IAEA-314-D4-RC-794.2
- Tuncbilek, A.S. (1997). Susceptibility of the saw-toothed grain beetle, *Oryzaephilus surinamensis* L. to gamma radiation. *Journal of Stored Products Research* 33 (4), 331-334.
- Tuncbilek, A.S., and I.A. Kansu (1996). The influence of rearing medium on the irradiation sensitivity of eggs and larvae of flour beetle, *Tribolium confusum* (J. du. Val). *Journal of Stored Products Research* 32 (1), 1-6.
- Tuncbilek, A.S. (1995). Effect of ⁶⁰Co gamma radiation on the rice weevil, *Sitophilus oryzae* (L.), *Anzeiger Fur Schadlingsskunde Pflanzenschutz Umweltschutz* 68 (2), 37-38.
- Yang, T.C.H., and H.S. Ducoff (1971). Recovery studies of X-irradiated *Tribolium castaneum* (Flour beetle) larvae. *Radiation Research* 46, 290-300.
- Wührer, B., and S.A. Hassan (1993). Selection of effective species/strains of *Trichogramma* (Hymenoptera: Trichogrammatidae) to control the diamondback moth *Plutella xylostella* L. (Lepidoptera: Plutellidae). *Journal of Applied Entomology* 116, 80-89.

