

ON TWO NEW CAVE SPECIES OF PSEUDOSCORPIONS (NEOBISIIDAE, PSEUDOSCORPIONES) FROM HERZEGOVINA AND DALMATIA

B. P. M. ĆURČIĆ¹, T. RAĐA², R. N. DIMITRIJEVIĆ¹, S. B. ĆURČIĆ¹, NINA B. ĆURČIĆ³ and
S. E. MAKAROV¹

¹ Institute of Zoology, Faculty of Biology, University of Belgrade, 11000 Belgrade, Serbia

² Speleological Society "Špiljar", 21000 Split, Croatia

³ Geographical Institute "Jovan Cvijić" SASA, 11000 Belgrade, Serbia

Abstract — A careful analysis of samples of pseudoscorpions (Neobisiidae, Pseudoscorpiones) from two underground habitats, one near Trebinje (Herzegovina) and the other from the National Park Paklenica (Croatia), has yielded two species of the genus *Roncus* L. Koch new to science, *Roncus tribunus* n. sp. and *R. argyrunti* n. sp. Both new species are described thoroughly, illustrated, and diagnosed. Some biogeographical and evolutionary characteristics of the two taxa are briefly discussed.

Key words: Pseudoscorpions, Neobisiidae, *Roncus tribunus* n. sp., *R. argyrunti* n. sp., caves, Herzegovina, Croatia.

INTRODUCTION


There exists no deeper karst than that situated in Dalmatia, Herzegovina and Montenegro. Not a drop of water remains long on the surface; all water sinks through various subterranean passages — potholes, ponors, fissures and crevices. Most water drains towards the Neretva River, while some drains towards the Zeta and Morača Rivers and Lake Skadar (Cvijić, 1926; Gavrilović, 1974). The study of pseudoscorpions in such underground habitats, a fauna whose composition is the result of a long and complicated history, enables us to evaluate the importance of the influence of long geomorphological isolation on the development of cave biota. Because of the age of these caves, different groups of pseudoscorpions have enjoyed the possibility of uninterrupted development and autochthonous differentiation. Such long isolation must have directed evolution towards greater

ecological differentiation within the framework of the cave system (Ćurčić, 1988).


In studying some Balkan pseudoscorpions from two small collections, we concentrated on two species belonging to the genus *Roncus* L. Koch (Neobisiidae). The former was represented by two females and one female, the latter by a single male. The specimens of *Roncus* from Herzegovina turned out to be a new species, *Roncus tribunus* n. sp., and a specimen of the genus from Croatia represents another new species — *Roncus argyrunti* n. sp. In this paper, both species are thoroughly described, diagnosed, and illustrated. In addition, some taxonomic, biogeographical, and evolutionary traits are briefly discussed.

MATERIALS AND METHODS

The pseudoscorpion specimens studied were mount-



Figs. 1 – 8. *Roncus tribunus* n. sp., holotype female from Herzegovina. 1 – pedipalpal chela, 2 – leg IV, 3 – pedipalp, 4 – epistome, 5 – carapace, 6 – flagellum, 7 – female genital area, 8 – chelicera. Scale lines = 0.25 mm (Figs. 1–3, and 5) and 0.50 mm (Figs. 4, 6, and 7).



Figs. 9 – 16. *Roncus argyruni* n. sp., holotype male from Croatia. 9 – pedipalpal chela, 10 – pedipalp, 11 – leg IV, 12 – epistome, 13 – carapace, 14 – flagellum, 15 – male genital area, 16 – chelicera. Scale lines = 0.25 mm (Figs. 9–11, and 13) and 0.50 mm (Figs. 12, 14–16).

ed on slides in gum chloral medium (Swan's fluid); they are deposited in the collections of the Institute of Zoology, Faculty of Biology, University of Belgrade, Belgrade, Serbia, and the Natural History Museum in Split, Croatia.

Setal designations follow Beier (1963).

SYSTEMATIC PART

NEOBISIIDAE J. C. CHAMBERLIN

RONCUS L. KOCH

RONCUS TRIBUNUS

B. ĆURČIĆ, NEW SPECIES
(Figs. 1-8; Table 1)

Etymology – After the Latin noun *Tribunus*, the primordial Roman name of Trebinje.

Material examined – Holotype female and paratype female from Kremni Do, Jama Bezdan Pit, nr. Trebinje, Herzegovina; 26 September 2008, collected by MP.

Description – Carapace considerably longer than broad (Fig. 5, Table 1). Neither eyes nor eyespots are developed (Fig. 5). Epistome small and low (Figs. 4 and 5). Carapacial setal formulae: $4 + 6 + 8 + 6 = 24$. Tergal formulae: $5 - 9 - 10 - 11 - 12 - 11 - 12 - 12 - 10 - 9$ and $6 - 7 - 9 - 11 - 10 - 12 - 12 - 12 - 10 - 9$. Female genital area: sternite II with 9-12 median posterior setae, sternite III with 14 or 15 posterior setae and four suprastigmatic microsetae on either side, sternite IV with 10 posterior setae and three or four small setae along each stigma (Fig. 7). Sternites V–X with $14 - 14 - 13 - 13 - 12 - 10$ and $14 - 14 - 13 - 15 - 14 - 11$ posterior setae. Twelfth abdominal segment with two pairs of small setae. Pleural membranes granulostrate.

Galea low and rounded (Fig. 8). Cheliceral palm with six setae, movable finger with one seta only (Fig. 8). Flagellum with one short proximal blade and six or seven longer blades distally, as is characteristic of

the genus *Roncus*. Dentition of cheliceral fingers as in Fig. 8.

Apex of pedipalpal coxa with five long acuminate setae. Small extero-lateral tubercle present on pedipalpal trochanter. Trochanter, femur and chelal palm with no granulations (Figs. 1 and 3). Fixed chelal finger with 61-69 teeth; movable chelal finger with 68-73 teeth. Chelal finger as long as chelal palm (Table 1). Disposition of trichobothria as illustrated in Fig. 1.

Tibia IV, metatarsus IV and tarsus IV each with a long sensitive seta (Fig. 2, Table 1).

Morphometric ratios and linear measurements (in mm) are presented in Table 1.

Distribution — Nr. Trebinje, Herzegovina, probably endemic (living in caves).

RONCUS ARGYRUNTI

B. ĆURČIĆ & RAĐA, NEW SPECIES
(Figs. 9-16; Table 1)

Etymology – After *Argyrunum*, an old Roman settlement near Starigrad, National Park Paklenica, Croatia.

Material examined – Holotype male from the Špilja Devnjača Cave, National Park Paklenica, Dalmatia, Croatia; 30 April 2001, collected by one of us (TR).

Description — Epistome small, low, triangular and apically rounded (Figs. 12 and 13). Neither eyes nor eyespots are developed (Figs. 12 and 13). Carapacial setal formula: $4 + 8 + 6 + 5 = 23$ setae. Tergite I – X setation: $6 - 9 - 10 - 11 - 12 - 11 - 11 - 11 - 10 - 10$. Male genital area: sternite II with 19 setae; sternite III with 7 anterior, 11 posterior and 3 suprastigmatic microsetae on either side; sternite IV with 10 posterior setae and 3 microsetae on either side; and sternites V–X each with $14 - 15 - 13 - 13 - 12 - 11$ posterior setae. Twelfth abdominal segment with two pairs of microsetae. Pleural membranes granulostrate.

Table 1. Linear measurements (in millimeters) and morphometric ratios in *Roncus tribunus* n. sp., and *R. argyrunti* n. sp. Abbreviations: FF = females, M = male.

Character	<i>R. tribunus</i> n. sp. FF	<i>R. argyrunti</i> n. sp. M
Body		
Length (1)	3.375-3.55	2.59
Cephalothorax		
Length (2)	0.93-0.94	0.72
Breadth (2a)	0.78-0.845	0.59
Ratio 2/2a	1.11-1.12	1.22
Abdomen		
Length	2.435-2.62	1.86
Chelicerae		
Length (3)	0.64-0.65	0.44
Breadth (4)	0.32-0.33	0.22
Length of movable finger (5)	0.46-0.47	0.295
Ratio 3/5	1.38-1.39	1.49
Ratio 3/4	1.94-2.03	2.00
Pedipalps		
Length with coxa (6)	5.50-5.835	3.61
Ratio 6/1	1.55-1.73	1.39
Length of coxa	0.77-0.78	0.56
Length of trochanter	0.61-0.64	0.45
Length of femur (7)	1.07-1.11	0.71
Breadth of femur (8)	0.295-0.305	0.22
Ratio 7/8	3.63-3.64	3.23
Ratio 7/2	1.14-1.19	0.99
Length of patella (tibia) (9)	0.94-0.95	0.61
Breadth of patella (tibia) (10)	0.39-0.40	0.275
Ratio 9/10	2.35-2.435	2.22
Length of chela (11)	2.03-2.05	1.28
Breadth of chela (12)	0.62	0.41
Ratio 11/12	3.31-3.33	3.12
Length of chelal palm (13)	1.01-1.05	0.63
Ratio 13/12	1.63-1.69	1.54
Length of chelal finger (14)	1.00-1.02	0.65
Ratio 14/13	0.95-1.01	1.03
Leg IV		
Total length	3.69-3.72	2.455
Length of coxa	0.51-0.52	0.37
Length of trochanter (15)	0.43-0.45	0.285
Breadth of trochanter (16)	0.19-0.20	0.13
Ratio 15/16	2.25-2.26	2.19
Length of femur + patella (17)	1.00-1.01	0.67
Breadth of femur + patella (18)	0.25-0.275	0.24
Ratio 17/18	3.67-4.00	2.77
Length of tibia (19)	0.91-0.93	0.58
Breadth of tibia (20)	0.14-0.15	0.13
Ratio 19/20	6.07-6.64	4.46
Length of metatarsus (21)	0.33-0.34	0.21
Breadth of metatarsus (22)	0.11	0.08
Ratio 21/22	3.00-3.09	2.625
Length of tarsus (23)	0.48-0.50	0.34
Breadth of tarsus (24)	0.10	0.07
Ratio 23/24	4.80-5.00	4.86
TS ratio - tibia IV	0.55-0.595	0.585
TS ratio - metatarsus IV	0.11-0.18	0.19
TS ratio - tarsus IV	0.36-0.37	0.36

Galea a low hyaline convexity (Fig. 16). Cheliceral palm with six setae, movable finger with one seta (Fig. 16). Flagellum with one short proximal blade and seven longer blades distally, characteristic of the genus (Fig. 14). Dentition of cheliceral fingers as in Fig. 16.

Manducatory process with four long setae. Small interior lateral tubercle present on pedipalpal trochanter (Fig. 10), femur and chelal palm with distinct interior granulations (Fig. 10), patella smooth. Group of microsetae proximal to trichobothria *eb* and *esb* not developed (Fig. 9); instead, four or five small setae distal to *eb* and *esb* present (Fig. 9). Fixed chelal finger with 56 teeth; movable chelal finger with 57 teeth. Chelal finger only slightly longer than chelal palm and shorter than pedipalpal femur (Table 1).

Trichobothriotaxy as illustrated in Figs. 9 and 10.

Tibia IV, metatarsus IV and tarsus IV each with a long sensitive seta (Fig. 11). Morphometric ratios and linear measurements (in mm) are presented in Table 1.

Distribution — Špilja Devnjača Cave, National Park Paklenica, Dalmatia, Croatia; probably an endemic and cave-dwelling form.

This study of pseudoscorpions inhabiting the Dinaric Karst offers further proofs of their great age and probably different origin. These species, or their stem forms, inhabited both leaf litter and humus of Dinaric forests during or even before the Tertiary. Finally, such forms represent the last vestiges of an old thermophilous and hygrophilous fauna, vestiges which found shelter in the underground domain (humus, soil, caves) of the Balkans and elsewhere (Čurčić, 1972, 1984, 1988, 1992a, b; Čurčić and Beron, 1981; Čurčić et al., 1993, 2004, 2010a, b, c, d, e, f, g; 2011a, b, c, d, e, f, g, h; 2012a, b, c, d, e, f; 2013a, b, c, d; Hadži, 1937).

Acknowledgment — Financial support for this work in the form of reimbursement for travel expenses was provided by the Serbian Ministry of Education, Science and Technological Development (Grant 173038).

REFERENCES

- Beier, M. (1963). Ordnung Pseudoscorpionidea (Afterskorpione). In: *Bestimmungsbücher zur Bodenfauna Europas*, Vol. 1. - Akademie Verlag, Berlin, 1-313.
- Cvijić, J. (1926). *Morphologie terrestre. II.* - Drž. štamp. Kralj. SHS, Belgrade. 1-506.
- Čurčić, B. P. M. (1972). Nouveaux pseudoscorpions cavernicoles de la Serbie et de la Macédoine. *Acta Mus. Mac. Sc. Nat. Skopje* **12**, 141-161.
- Čurčić, B. P. M. (1984). On two new species of *Roncus* L. Koch, 1873, from Macedonia (Arachnida: Pseudoscorpiones: Neobisiidae). *Senckenbergiana biol.*, **65**, 97-104.
- Čurčić, B. P. M. (1988). Cave-Dwelling Pseudoscorpions of the Dinaric Karst. - *Acad. Sci. Art. Slov., Cl. IV, Hist. Nat., Opera* **26**, *Inst. Biol. Ioannis Hadži*, **8**, Ljubljana, 1-192.
- Čurčić, B. P. M. (1992a). New and little-known pseudoscorpions of the genus *Roncus* L. Koch (Neobisiidae, Pseudoscorpiones) from Serbia, Yugoslavia. *Bijdr. Dierk.*, **61** (4), 237-249.
- Čurčić, B. P. M. (1992b). A new species of *Roncus* L. Koch, 1873, from southeastern Europe (Pseudoscorpiones: Neobisiidae). *Proc. Entomol. Soc. Wash.*, **94** (4), 447-453.
- Čurčić, B. P. M. and P. Beron (1981). New and little-known cavernicole pseudoscorpions in Bulgaria (Neobisiidae, Pseudoscorpiones, Arachnida). *Glasn. Srpsk. Akad. Nauka, Beograd*, 329; *Cl. Sci. Nat.-Math.*, **48**, 63-85.
- Čurčić, B. P. M., Čurčić, S. B., Čurčić, N. B. and S. E. Makarov (1993). Three new epigean representatives of *Roncus* L. Koch, 1873 (Neobisiidae, Pseudoscorpiones), from the Balkan Peninsula. *Bijdr. Dierk.*, **62** (4), 237-248.
- Čurčić, B. P. M., Dimitrijević, R. N. and A. Legakis (2004). The Pseudoscorpions of Serbia, Montenegro, and the Republic of Macedonia. Monographs, 8. - Institute of Zoology - Faculty of Biology - University of Belgrade, Hellenic Zoological Society, Committee for Karst and Speleology - Serbian Academy of Sciences and Arts, Institute of Nature Conservation of the Republic of Serbia, Belgrade-Athens, 1-400.
- Čurčić, B. P. M., Dimitrijević, R. N. and N. B. Čurčić (2010a). *Neobisium deltshevi* (Neobisiidae, Pseudoscorpiones), a new endemic cave-dwelling pseudoscorpion from East Serbia. *Arch. Biol. Sci., Belgrade*, **62** (1), 191-198.

- Ćurčić, B. P. M., Dimitrijević, R. N., Rađa, T., Ćurčić, N. B. and M. Milinčić (2010b). *Chthonius* (*Chthonius*) *onaei* n. sp. (Chthoniidae, Pseudoscorpiones), a new epigean species from Croatia. *Arch. Biol. Sci., Belgrade*, **62** (2), 494-499.
- Ćurčić, B. P. M., Rađa, T., Ćurčić, S. B. and N. B. Ćurčić (2010c). On *Roncus almissae* n. sp., *R. krupanjensis* n. sp., and *R. radji* n. sp., three new pseudoscorpions (Pseudoscorpiones, Neobisiidae) from Croatia and Serbia, respectively. *Arch. Biol. Sci., Belgrade*, **62** (2), 503-513.
- Ćurčić, B. P. M., Makarov, S. E., Rađa, T., Ćurčić, S. B., Ćurčić, N. B. and M. Pecelj (2010d). On three new cave pseudoscorpions (Pseudoscorpiones, Neobisiidae) from Mt. Mosor, Dalmatia (Croatia). *Arch. Biol. Sci., Belgrade*, **62** (3), 813-828.
- Ćurčić, B. P. M., Lemaire, J.-M., Ćurčić, S. B., Dimitrijević, R. N., Milinčić, M. and M. Pecelj (2010e). Two new epigean pseudoscorpions (Neobisiidae, Pseudoscorpiones) from the Maritime Alps, France. *Arch. Biol. Sci., Belgrade*, **62** (3), 829-834.
- Ćurčić, B. P. M., Rađa, T., Dimitrijević, R. N., Tomić, V. T., Pecelj, M. and S. B. Ćurčić (2010f). *Chthonius* (*Chthonius*) *torakensis* and *Chthonius* (*Ephippiochthonius*) *cikolae*, two new species of pseudoscorpions (Chthoniidae) from Croatia. *Arch. Biol. Sci., Belgrade*, **62** (4), 1223-1229.
- Ćurčić, B. P. M., Rađa, T., Dimitrijević, R. N., Ćurčić, S. B. and M. Milinčić (2010g). A new troglomorphic pseudoscorpion (Pseudoscorpiones, Neobisiidae) from Istria, Croatia. *Arch. Biol. Sci., Belgrade*, **62** (4), 1245-1250.
- Ćurčić, B. P. M., Dimitrijević, R. N., Makarov, S. E., Milinčić, M., Pecelj, M. and T. Rađa (2011a). Two new pseudoscorpions from the UN Administered Province of Kosovo and Croatia. *Arch. Biol. Sci., Belgrade*, **63** (1), 235-244.
- Ćurčić, B. P. M., Ćurčić, S. B., Ćurčić, N. B. and B. S. Ilić (2011b). *Chthonius* (*Globochthonius*) *medeonis* n. sp., a new cave false scorpion from Montenegro. *Arch. Biol. Sci., Belgrade*, **63** (1), 245-250.
- Ćurčić, B. P. M., Rađa, T., Makarov, S. E., Ćurčić, S. B., Ilić, B. S. and R. N. Dimitrijević (2011c). A cavernicolous pseudoscorpion of the genus *Chthonius* (*Chthonius*) C. L. Koch from Dalmatia. *Arch. Biol. Sci., Belgrade*, **63** (2), 493-498.
- Ćurčić, B. P. M., Dimitrijević, R. N. and N. B. Ćurčić (2011d). A new cave pseudoscorpion (Pseudoscorpiones: Chthoniidae): *Chthonius* (*Chthonius*) *lupinus* n. sp. from Bosnia-Herzegovina. *Arch. Biol. Sci., Belgrade*, **63** (2), 499-506.
- Ćurčić, B. P. M., Ilić, B. S., Makarov, S. E. and V. T. Tomić (2011e). On two new cave-dwelling and relict pseudoscorpions of the genus *Chthonius* C. L. Koch (Chthoniidae, Pseudoscorpiones) from Bosnia. *Arch. Biol. Sci., Belgrade*, **63** (3), 847-854.
- Ćurčić, B. P. M., Ćurčić, S. B., Ćurčić, N. B., Rađa, T. and R. N. Dimitrijević (2011f). On two new pseudoscorpions from Herzegovina. *Arch. Biol. Sci., Belgrade*, **63** (3), 855-865.
- Ćurčić, B. P. M., Rađa, T., Dimitrijević, R. N., Ćurčić, N. B., Ilić, B. S. and J. M. Pecelj (2011g). *Chthonius* (*Ephippiochthonius*) *lagadini* n. sp. (Chthoniidae, Pseudoscorpiones), a new endemic epigean pseudoscorpion from Macedonia. *Arch. Biol. Sci., Belgrade*, **63** (4), 1251-1256.
- Ćurčić, B. P. M., Rađa, T., Dimitrijević, R. N., Makarov, S. E. and M. Milinčić (2011h). A new cave pseudoscorpion from Serbia (Pseudoscorpiones, Chthoniidae). *Arch. Biol. Sci., Belgrade*, **63** (4), 1257-1263.
- Ćurčić, B. P. M., Stojanović, D. Z., Ilić, B. S. and N. B. Ćurčić (2012a). *Roncus ivansticae* (Neobisiidae, Pseudoscorpiones), a new epigean species from Eastern Serbia. *Arch. Biol. Sci., Belgrade*, **64** (1), 371-377.
- Ćurčić, B. P. M., Makarov, S. E., Ćurčić, S. B., Antić, D. Ž. and R. N. Dimitrijević (2012b). A new soil pseudoscorpion, *Roncus ursi* n. sp., from Western Serbia (Neobisiidae, Pseudoscorpiones). *Arch. Biol. Sci., Belgrade*, **64** (1), 379-384.
- Ćurčić, B. P. M., Dimitrijević, R. N., Tomić, V. T., Pecelj, M., Ilić, B. S., Makarov, S. E., Ćurčić, N. B. and S. Pešić (2012c). A new epigean false scorpion: *Roncus sumadijiae* n. sp. (Neobisiidae, Pseudoscorpiones) from the Balkan Peninsula (Western Serbia). *Arch. Biol. Sci., Belgrade*, **64** (2), 703-708.
- Ćurčić, B. P. M., Rađa, T. and R. N. Dimitrijević (2012d). On two new cave pseudoscorpions, *Chthonius* (*Chthonius*) *pagus* n. sp. (Chthoniidae) and *Roncus navalia* (Neobisiidae), from the Island of Pag, Croatia. *Arch. Biol. Sci., Belgrade*, **64** (4), 1555-1565.
- Ćurčić, B. P. M., Makarov, S. E., Rađa, T., Ilić, B. S. and D. Ž. Antić (2012e). *Roncus meledae* n. sp. and *Neobisium oculatum* n. sp., from the Island of Mljet, Dalmatia (Neobisiidae, Pseudoscorpiones). *Arch. Biol. Sci., Belgrade*, **64** (4), 1567-1576.
- Ćurčić, B. P. M., Dimitrijević, R. N., Rađa, T., Makarov, S. E. and B. S. Ilić (2012f). *Archaeoroncus*, a new genus of pseudoscorpions from Croatia (Pseudoscorpiones, Neobisiidae), with descriptions of two new species. *Acta Zool. Bulg.*, **64** (4), 333-340.
- Ćurčić, B. P. M., Dimitrijević, R. N., Makarov, S. E., Ćurčić, S. B., Tomić, V. T., Antić, D. Ž., Ilić, B. S. and N. B. Ćurčić (2013a). *Roncus radgost* n. sp., *R. jarevid* n. sp., and *R. crnobog* n. sp., three new cave-dwellers from Eastern Serbia. *Arch. Biol. Sci., Belgrade*, **65** (2), 751-760.
- Ćurčić, B. P. M., Rađa, T., Makarov, S. E., Dimitrijević, R. N., Ćurčić, S. B. and B. S. Ilić (2013b). On the identity of types of *Roncus diocletiani* Ćurčić, Dimitrijević & Rađa and *Archaeoroncus tenuis* (Hadži) (Neobisiidae, Pseudoscor-

- piones) from Croatia. *Arch. Biol. Sci., Belgrade*, **65** (2), 761-766.
- Ćurčić, B. P. M., Dimitrijević, R. N., Tomić, V. T., Makarov, S. E., Ćurčić, N. B. and S. B. Ćurčić (2013c). Biodiversity of *Roncus* L. Koch in Montenegro – *Roncus teutae* n. sp. from Mt. Orjen (Neobisiidae, Pseudoscorpiones). *Arch. Biol. Sci., Belgrade*, **66** (1), 369-375.
- Ćurčić, B. P. M., Rađa, T., Dimitrijević, R. N., Ćurčić, S. B., Makarov, S. E. and D. Ž. Antić (2013d). A new pseudoscorpion from Bosnia: *Roncus bosniensis* n. sp. (Neobisiidae, Pseudoscorpiones). *Arch. Biol. Sci., Belgrade*, **66** (1), 363-368.
- Gavrilović, D. (1974). Die serbische Karstterminologie. - *Verh. geogr. Inst. Jugosl., Komm. Karstterm. jugosl. Volker*, 2, Belgrade. 1-73.
- Hadži, J. (1937). Pseudoskorpioniden aus Südserbien. *Glasnik Skopskog naučnog društva, Skopje*, **18**, 13-38.